



King's Research Portal

DOI:

[10.1159/000486546](https://doi.org/10.1159/000486546)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Knight, R., Khondoker, M., Magill, N., Stewart, R., & Landau, S. (2018). A systematic review and meta-analysis of the effectiveness of acetylcholinesterase inhibitors and memantine in treating the cognitive symptoms of dementia. *Dementia and Geriatric Cognitive Disorders*, 45, 131–151. <https://doi.org/10.1159/000486546>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

A systematic review and meta-analysis of the effectiveness of acetylcholinesterase inhibitors and memantine in treating the cognitive symptoms of dementia

Ruth Knight^{a*}, Mizanur Khondoker^b, Nicholas Magill^a, Robert Stewart^{c,d}, Sabine Landau^a

a Department of Biostatistics and Health Informatics, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK

b Norwich Medical School, University of East Anglia, Norwich, UK

c Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King's College London

d South London and Maudsley NHS Foundation Trust, London, UK

* Corresponding author: Department of Biostatistics and Health Informatics, Institute of Psychiatry, Psychology and Neuroscience, King's College London, De Crespigny Park, London, SE5 8AF, UK; +44 (0)20 7848 0983; ruth.s.young@kcl.ac.uk

Short title: AChEIs and memantine in treating the cognitive symptoms of dementia

Conflicts of interest: None

Funding: This paper represents independent research funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

Keywords: Acetylcholinesterase inhibitor, memantine, AD, Vascular dementia, systematic review, meta-analysis

Abstract

Background: Acetylcholinesterase inhibitors (AChEIs) and memantine are commonly used in the management of dementia. In routine clinical practice, dementia is often monitored via the mini-mental state examination (MMSE). We conducted a systematic review and meta-analysis of the effects of these drugs on MMSE scores. **Summary:** Eighty trials were identified. Pooled effect estimates were in favour of both AChEIs and memantine at 6 months. Meta-regression indicated that dementia sub-type was a moderator of AChEI treatment effect with the effect of treatment versus control twice as high for PDD/DLB patients (2.11 MMSE points at 6 months) as for AD/VaD patients (0.91 MMSE points at 6 months). **Key messages:** AChEIs demonstrate a modest effect versus control on MMSE scores which is moderated by dementia sub-type. For memantine the effect is smaller.

Introduction

Dementia is a major health concern in elderly populations worldwide which can affect many aspects of a person's life and functioning. There is currently no cure for most forms of dementia but several drugs are used in its management. The acetylcholinesterase inhibitors (AChEIs) were developed as a consequence of the cholinergic hypothesis of cognitive decline [1] and the NMDA receptor agonist memantine as a consequence of an hypothesised role of the glutamatergic system in neurodegeneration [2]. The effectiveness of these treatments has been evaluated in a large number of randomised controlled trials (RCTs) across functional, global, cognitive and neuropsychiatric domains [3-5]. This review focuses on their effects on cognition.

Measures of global cognition include the mini-mental state examination (MMSE) [6], the Alzheimer's disease assessment scale - cognitive subscale (ADAS-cog) [7], and the Severe Impairment Battery (SIB) [8], which focuses on those with severe cognitive impairment. Existing meta-analyses tend either to consider cognitive outcomes on the ADAS-cog or SIB [9] or to use standardised mean differences to combine results from several scales [10]. In this review results are analysed relating to the MMSE scale specifically. A small number of existing meta-analyses combine cognitive outcomes on the MMSE; however, these are mainly focused on diagnostic and medication subgroups and do not cover all available trials. The largest of these includes only 21 MMSE effect estimates [11], less than half of the number included in this review.

The MMSE is the scale most often used to monitor dementia severity and progression in routine clinical practice, and thus the advantage of reviewed outcomes on this scale is better clinical interpretability and relevance to routine care. In addition, the volume of evidence can be substantially increased by the inclusion of ADAS-cog results translated to MMSE scale equivalents.

Methods

A protocol for this systematic review was prospectively registered on PROSPERO and can be found at https://www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42015025892.

Search strategy

A two-tier search strategy was employed to identify relevant trials for inclusion in this review. First, existing systematic reviews and meta-analyses assessing the drugs of interest were identified and citations to included trials extracted. Following this, additional searches subdivided by dementia diagnosis and, where necessary, drug received, were conducted to identify trials published since the date of the most recent review.

Searches were conducted using the Web of Science, MEDLINE, PsycINFO, EMBASE and CINAHL databases. Final searches were conducted in March 2017. Searches were combinations of: (i) drug names e.g. “donepezil”, “galantamine”, “rivastigmine”, “memantine”; (ii) diagnoses e.g. “Alzheimer*”, “vascular dement*”, “lewy* bod*”, “Parkinson* disease dement*”; and (iii) “randomi?ed” and “trial”. A full list of search terms used is provided in the supplementary material. Further searches were carried out using the International Clinical Trials Registry Platform (ICTRP) and industry trial registers to identify unpublished trials. References and citing articles of selected trials were assessed to identify further trials for inclusion.

Study selection criteria and data extracted

Trials were included if they met the following criteria: (i) a randomised trial designed to evaluate the effectiveness of AChEI monotherapy, memantine monotherapy or memantine treatment in a group of patients some, but not all, of whom received a concurrent AChEI; (ii) treatments compared to a control group receiving placebo or no treatment; (iii) participants in the trial diagnosed with Alzheimer’s disease (AD), vascular dementia (VaD), Parkinson’s disease dementia (PDD), dementia with Lewy bodies (DLB) or frontotemporal dementia (FTD); (iv) at least one of the MMSE or ADAS-cog used as an outcome; and (v) sufficient data provided, defined as at least one treatment effect estimate and associated standard error (SE) on either the MMSE or ADAS-cog. Treatment effect estimates used included change score differences and time point differences. In some cases, effect estimates and SEs had to be calculated from other statistics (for example, confidence intervals).

From each trial, data were extracted on: (i) Trial design – duration, inclusion and exclusion criteria, numbers of patients randomised to each arm, intervention and control conditions, type of randomisation, details on blinding, cognitive assessments and measurement times; (ii) Analysis approaches – analysis method, missing data methods and effect size estimate used; and (iii) Trial data – baseline data, attrition and adherence rates, treatment effect estimates and SEs.

Study selection and data extraction were conducted by one reviewer (RK) and a sample of each was checked by a second reviewer (NM). Reviewers agreed on study selection in 99% of cases and agreement regarding data extraction was also high: 87.5% for risk of bias assessment, 82.8% for baseline measures and 75% for effect estimates. Most effect estimate discrepancies were due to miscommunication on how these were extracted. All discrepancies were discussed and resolved.

ADAS-cog translation

The objective of the meta-analysis was to estimate the treatment effect on the MMSE; however, effect estimates on the ADAS-cog were also collected and translated, since both scales measure global cognition. Baseline measures from the 36 trials which measured both were used to translate. MMSE scores range from 0 to 30 and ADAS-cog scores from 0 to 70 and both MMSE=30 and ADAS-cog=0 represent healthy cognition. Thus, a linear regression of ADAS-cog on MMSE with intercept fixed at 30 was fitted. The resulting model was: $MMSE = 30 - 0.42 * ADAS-cog$, with a squared multiple correlation of 0.679 suggesting fairly good fit. Translation of both treatment effect estimates and SEs required only the coefficient. Treatment effect estimates were translated using $MMSE = -0.42 * ADAS-cog$, and the SEs using $MMSE = 0.42 * ADAS-cog$.

Risk of bias assessment

The risk of bias in included studies was assessed using the Cochrane risk of bias tool [12]. This determines whether the risk of internal bias under a series of domains is low, high or unclear. These were combined so that a trial rated low in all domains was at low risk of bias. One domain, reporting bias, was excluded from the combination, since trial protocols were required to assess it but were not available for most included trials due to their age.

Statistical analyses

Random-effects meta-analysis [13] was used to combine trial results. This was conducted separately for AChEIs and memantine. Pooled effects were estimated at 3, 6 and 12 months (± 14 days) after treatment initiation. Effect estimates were also considered in AChEI drug subgroups. Heterogeneity was assessed using the I^2 statistic [14] and publication bias using funnel plots and Begg and Mazumdar's rank correlation test [15]. All statistical analyses were conducted using R [16] and the metafor package [17].

Meta-regressions were conducted to assess the impact of data quality on effect size estimates and test potential moderators. The data quality factors were: (i) the inclusion of translated results; and (ii) the risk of bias assessment overall rating. The hypothesised potential moderators were: (i) AChEI (donepezil, galantamine or rivastigmine); (ii) dementia diagnosis (AD, VaD, PDD/DLB or FTD); (iii) baseline MMSE score; and (iv) date of publication (before or after 2000). All were categorical factors except baseline MMSE, which was continuous. The Knapp and Hartung adjustment [18] was used to account for uncertainty in the assessment of residual heterogeneity. The omnibus test of coefficients was used to identify factors significant at the 5% and 1% levels.

Results

Literature search results

The search for systematic reviews identified 522 citations of which 52 were relevant, and these included 194 citations to trials. An additional 857 citations were identified by further searches for trials, resulting in 1051 possible citations. After removal of duplicates, title and abstract screening, and full text screening, 84 references about 74 trials met the inclusion criteria. Searches in ICTRP and industry

registers and citation tracking identified a further 6 trials for inclusion. In total, 80 trials met the inclusion criteria. The process of identifying these is detailed in Figure 1.

Characteristics of included studies

Of the included trials summarised in Table 1, half (40) investigated donepezil and the others were evenly split amongst galantamine (13), rivastigmine (14) and memantine (13). The majority of the trials (55) were conducted in patients with AD. Other diagnoses were VaD (9), AD and VaD (4), PDD or DLB (10) and FTD (2). Dementia severity ranged from mild in some trials to severe in others. The trials lasted between 4 and 104 weeks and many recorded outcome measures at intermediate time points. Forty-eight trials provided MMSE outcomes, twenty-four ADAS-cog and the remainder reported a mixture of the two.

The average baseline age in AChEI trials was 73.8 years and in memantine trials was 75.9 years. The proportion of women was slightly more than half in the AChEI trials (mean 57.5%; range 7.1%-84.6%), and the memantine trials (mean 56.3%; range 25%-73.8%). The mean baseline MMSE was higher in the AChEI trials (18.6 points) than in the memantine trials (16.5).

Risk of bias assessment

The Cochrane risk of bias tool was applied to each trial and the final column of Table 1 records overall ratings. Risk of bias was low in 14 trials, high in 45 trials and unclear in 21 trials. The large number of trials rated high risk was mainly due to missing data methods combined with relatively high volumes of missing data. The majority of trials used observed case or last observation carried forward analyses which both introduce a significant risk of bias in the presence of missing data.

Meta-analysis results

AChEIs – 3 months

At 3 months (± 14 days) after treatment initiation, 42 trials provided 60 estimates of treatment effect. The pooled effect estimate (Figure 2) was 1.08 MMSE points (95% CI 0.92-1.23). There was evidence of heterogeneity ($I^2=68.2\%$) and this was later explored via meta-regression. Begg and Mazumdar's rank test suggested some publication bias ($p=0.01$) and the funnel plot supported this (Figure 3); however, the patterns did not seem overly concerning. In the drug subgroups the treatment effects ranged from 0.98 (95% CI 0.32-1.63) for rivastigmine to 1.15 (95% CI 0.69-1.61) for donepezil 3-5mg/d.

AChEIs – 6 months

At 6 months (± 14 days) after treatment initiation, 38 trials provided 52 estimates of treatment effect. The pooled effect estimate was 1.00 (95% CI 0.83-1.16; Figure 4), and there was evidence of heterogeneity ($I^2=69.9\%$). Neither the funnel plot nor the rank correlation test ($p=0.385$) suggested publication bias. The effect estimates in treatment subgroups ranged from 0.69 (95% CI 0.43-0.95) for rivastigmine to 1.39 (95% CI 0.79-2.00) for galantamine.

AChEIs – 12 months

At 12 months (± 14 days) after treatment initiation, 4 trials provided estimates of treatment effect. The pooled effect estimate was 1.10 (95% CI 0.48-1.72; Figure 5). There was evidence for heterogeneity ($I^2=79\%$); however, the funnel plot did not suggest any obvious publication bias and there were too few estimates for a formal test.

Memantine – 3, 6 and 12 months

Treatment effect estimates were provided by 12 memantine trials: 4 at 3 months; 8 at 6 months; and 3 at 12 months after treatment initiation. The pooled effect estimates at each time point were in favour of treatment though were much smaller than those for the AChEIs (Figure 6). At 12 months the pooled effect did not reach significance (0.41, 95% CI -0.44 to 1.26). At all 3 time points the I^2 values were small suggesting little heterogeneity.

Meta-regressions

High I^2 values observed for the AChEI meta-analyses at 3 and 6 months suggested considerable variability in the effect estimates; this was investigated further via meta-regression. Factors investigated were data quality measures and potential moderators, as listed in the methods section. Tables 2 and 3 provide meta-regression coefficients, associated p-values and the p-value for the omnibus test of parameters at 3 and 6 months respectively. For categorical factors, coefficients are the difference in average effect estimates for each category versus the reference category; for continuous factors, they are the relation between the factor and the effect estimate. Factors for which the omnibus test of parameters was significant at the 5% and 1% levels are highlighted.

A true moderator of treatment effect would be expected to last over time, thus, only factors significant at both 3 and 6 months were considered. Dementia sub-type diagnosis was the only factor significant at both 3 months ($p=0.009$) and 6 months ($p=0.007$). Examination of diagnostic subgroup results suggested that the effects in the AD and VaD subgroups were the same but those in the PDD/DLB subgroup were different.

Meta-analyses in diagnosis subgroups

At 3 months the pooled effect estimate in the AD/VaD subgroup was 0.97 MMSE points (95% CI 0.85-1.10) and in the PDD/DLB subgroup 1.99 (1.18-2.81). At 6 months the effect in the AD/VaD subgroup was 0.91 (0.77-1.05) and in the PDD/DLB subgroup was 2.11 (0.61-3.61). All four trials providing an effect estimate at 12 months were in the AD/VaD subgroup. The memantine trials provided too few trials for meta-regression to be conducted; however, at both 6 months and 12 months the effects in the PDD/DLB subgroup were significantly higher (1.90 points at 6 months and 1.80 points at 12 months) than those in the AD/VaD subgroup (0.36 points at 6 months and 0.31 points at 12 months).

Discussion

This review identified 80 trials evaluating the effects of donepezil, galantamine, rivastigmine and memantine on cognitive function in dementia, more than in any previous review. Cognitive effects were extracted on the MMSE, the outcome of interest, or the ADAS-cog. Baseline measures from 36 trials which measured both were used to enable translation of ADAS-cog results to the MMSE scale. This allowed the inclusion of 24 additional trials and results at additional time points from a further 8 trials. The large number of studies included in this review is one of its strengths and this number is increased through the translation of ADAS-cog results. The translation relationship has good R^2 ; however, this relationship has not been used elsewhere and should therefore be treated as preliminary and requiring confirmation.

Meta-regressions of the AChEI results at 3 and 6 months identified one moderator of treatment effect, dementia sub-type diagnosis. Treatment effects were smaller for those patients diagnosed with AD or VaD (0.97 MMSE points at 3 months and 0.91 points at 6 months) than those diagnosed with PDD or DLB (1.99 points at 3 months and 2.11 points at 6 months). All trials reporting effects at 12 months were for AD or VaD patients and these indicated a similar effect to those at 3 and 6 months (1.10 points). The higher response seen in the PDD/DLB group is consistent with previous results [19] and may be due to the greater cholinergic deficit seen in these conditions [20]. The effects observed in the AD/VaD subgroup are somewhat smaller than those in a previous review of AChEIs in AD only [5]. This may be due to the inclusion of VaD results, which evidence suggests may give rise to more mixed findings on AChEI effect [21,22]; however, meta-regression indicated no significant differences between AD and VaD subgroups. Whilst these drugs are only licensed for the use in AD or PDD, there is evidence that they are widely used for patients with DLB and VaD in routine clinical practice [23], and thus the inclusion of these trial results was felt to be appropriate.

The number of trials providing estimates of memantine treatment effects was much smaller and it was not possible to conduct meta-regression analyses; however, results were calculated for the previously identified subgroups. In the AD/VaD subgroup the effects were small and in favour of treatment (0.65 MMSE points at 3 months, 0.36 points at 6 months and 0.31 points at 12 months). Again, the effects in the PDD/DLB subgroup were higher (1.90 points at 6 months and 1.80 points at 12 months). Few of these effects were significantly different from zero.

Through the results of this review, we sought to increase clinical interpretability and relevance to routine care since they are estimated on the MMSE, the scale most often used to monitor dementia in clinical practice. Estimation of MMSE effects also potentially enables results to be compared, contrasted and in future combined with observational findings from routine clinical practice. The AChEI results suggest a treatment effect of around one MMSE point at 3, 6 and 12 months after treatment initiation. Since studies have suggested that the annual rate of MMSE decline amongst dementia patients is 4 to 5 MMSE points [24], such an effect estimate is modest: equivalent to an approximately 3 month delay in cognitive decline. However, while the effect sizes are small, they could have a significant impact in terms of costs and hospital or nursing home admissions, which have both been shown to be linked to level of cognitive function as measured by the MMSE score [25]. In addition, the length of time for which these benefits continue may be of interest [23].

Use of the MMSE scale makes the results of this review more clinically applicable; however, there are several limitations to this scale. It suffers from both floor and ceiling effects [26], though these should not be of particular concern for the trials included in this study. In addition, it is particularly suitable for measuring the cognitive deficits observed in AD and may be less sensitive to those in VaD [27] or FTD [28]. However, the latter has little impact in the current review since only one included trial concerned FTD and, as mentioned, no significant differences were found between AD and VaD sub-groups in meta-regressions.

References

- 1 Terry AV, Buccafusco J: The cholinergic hypothesis of age and alzheimer's disease-related cognitive deficits: Recent challenges and their implications for novel drug development. *Journal of Pharmacology and Experimental Therapeutics* 2003;306:821-827.
- 2 Cacabelos R, Takeda M, Winblad B: The glutamatergic system and neurodegeneration in dementia: Preventive strategies in alzheimer's disease. *International journal of geriatric psychiatry* 1999;14:3-47.
- 3 Van De Glind EMM, Van Enst WA, Van Munster BC, Olde Rikkert MGM, Scheltens P, Scholten RJP, Hooft L: Pharmacological treatment of dementia: A scoping review of systematic reviews. *Dement Geriatr Cogn Disord* 2013;36:211-228.
- 4 Winblad B, Jones RW, Wirth Y, Stoffler A, Mobius HJ: Memantine in moderate to severe alzheimer's disease: A meta-analysis of randomised clinical trials. *Dement Geriatr Cogn Disord* 2007;24:20-27.
- 5 Birks J: Cholinesterase inhibitors for alzheimer's disease. *Cochrane Database of Systematic Reviews* 2006;(1)
- 6 Folstein MF, Folstein SE, McHugh PR: "Mini-mental state": A practical method for grading the cognitive state of patients for the clinician. *Journal of psychiatric research* 1975;12:189-198.
- 7 Rosen WG, Mohs RC, Davis KL: A new rating scale for alzheimer's disease. *The American journal of psychiatry* 1984
- 8 Panisset M, Roudier M, Saxton J, Boiler F: Severe impairment battery: A neuropsychological test for severely demented patients. *Archives of neurology* 1994;51:41-45.
- 9 Tan C-C, Yu J-T, Wang H-F, Tan M-S, Meng X-F, Wang C, Jiang T, Zhu X-C, Tan L: Efficacy and safety of donepezil, galantamine, rivastigmine, and memantine for the treatment of alzheimer's disease: A systematic review and meta-analysis. *Journal of Alzheimer's Disease* 2014;41:615-631.
- 10 Di Santo SG, Prinelli F, Adorni F, Caltagirone C, Musicco M: A meta-analysis of the efficacy of donepezil, rivastigmine, galantamine, and memantine in relation to severity of alzheimer's disease. *Journal of Alzheimer's Disease* 2013;35:349-361.
- 11 Raina P, Santaguida P, Ismail A, Patterson C, Cowan D, Levine M, Booker L, Oremus M: Effectiveness of cholinesterase inhibitors and memantine for treating dementia: Evidence review for a clinical practice guideline. *Ann Intern Med* 2008;148:379-397.
- 12 Higgins J, Green S: *Cochrane handbook for systematic reviews of interventions*. Version 5.1. 0. The cochrane collaboration; 2011, 2013,
- 13 DerSimonian R, Laird N: Meta-analysis in clinical trials. *Controlled clinical trials* 1986;7:177-188.
- 14 Higgins J, Thompson SG: Quantifying heterogeneity in a meta-analysis. *Statistics in medicine* 2002;21:1539-1558.
- 15 Begg CB, Mazumdar M: Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994;1088-1101.
- 16 R Core Team: *R: A language and environment for statistical computing*. R foundation for statistical computing, vienna, austria, 2014,
- 17 Viechtbauer W: Conducting meta-analyses in r with the metafor package. *J Stat Softw* 2010;36:1-48.
- 18 Knapp G, Hartung J: Improved tests for a random effects meta-regression with a single covariate. *Statistics in medicine* 2003;22:2693-2710.
- 19 Samuel W, Caligiuri M, Galasko D, Lacro J, Marini M, McClure FS, Warren K, Jeste DV: Better cognitive and psychopathologic response to donepezil in patients prospectively diagnosed as dementia with lewy bodies: A preliminary study. *International journal of geriatric psychiatry* 2000;15:794-802.

- 20 Tiraboschi P, Hansen L, Alford M, Sabbagh M, Schoos B, Masliah E, Thal L, Corey-Bloom J: Cholinergic dysfunction in diseases with lewy bodies. *Neurology* 2000;54:407-407.
- 21 Birks J, McGuinness B, Craig D: Rivastigmine for vascular cognitive impairment. *Cochrane Database of Systematic Reviews* 2013
- 22 Birks J, Harvey RJ: Donepezil for dementia due to alzheimer's disease. *The Cochrane Library* 2006
- 23 Perera G, Khondoker M, Broadbent M, Breen G, Stewart R: Factors associated with response to acetylcholinesterase inhibition in dementia: A cohort study from a secondary mental health care case register in london. *PLoS One* 2014;9:e109484.
- 24 Ballard C, O'brien J, Morris C, Barber R, Swann A, Neill D, McKeith I: The progression of cognitive impairment in dementia with lewy bodies, vascular dementia and alzheimer's disease. *International journal of geriatric psychiatry* 2001;16:499-503.
- 25 Knapp M, Chua K-C, Broadbent M, Chang C-K, Fernandez J-L, Milea D, Romeo R, Lovestone S, Spencer M, Thompson G: Predictors of care home and hospital admissions and their costs for older people with alzheimer's disease: Findings from a large london case register. *BMJ open* 2016;6:e013591.
- 26 Franco-Marina F, García-González JJ, Wagner-Echeagaray F, Gallo J, Ugalde O, Sánchez-García S, Espinel-Bermúdez C, Juárez-Cedillo T, Rodríguez MÁV, García-Peña C: The mini-mental state examination revisited: Ceiling and floor effects after score adjustment for educational level in an aging mexican population. *International psychogeriatrics* 2010;22:72-81.
- 27 Dong Y, Sharma VK, Chan BP-L, Venketasubramanian N, Teoh HL, Seet RCS, Tanicala S, Chan YH, Chen C: The montreal cognitive assessment (moca) is superior to the mini-mental state examination (mmse) for the detection of vascular cognitive impairment after acute stroke. *Journal of the neurological sciences* 2010;299:15-18.
- 28 Osher JE, Wicklund AH, Rademaker A, Johnson N, Weintraub S: The mini-mental state examination in behavioral variant frontotemporal dementia and primary progressive aphasia. *American Journal of Alzheimer's Disease & Other Dementias®* 2008;22:468-473.
- 29 Frolich L, Ashwood T, Nilsson J, Eckerwall G, Sirocco I: Effects of azd3480 on cognition in patients with mild-to-moderate alzheimer's disease: A phase iib dose-finding study. *Journal of Alzheimers Disease* 2011;24:363-374.
- 30 Gault LM, Ritchie CW, Robieson WZ, Pritchett Y, Othman AA, Lenz RA: A phase 2 randomized, controlled trial of the alpha7 agonist abt-126 in mild-to-moderate alzheimer's dementia. *Alzheimer's and Dementia: Translational Research and Clinical Interventions* 2015;1:81-90.
- 31 Geldmacher DS, Perdomo C, Pratt R: Effect of donepezil treatment on visual attention/exploration tasks in patients with mild to moderately severe alzheimer's disease: Results of a pilot study. *Neurobiology of Aging* 2000;21:S169.
- 32 Marek GJ, Katz DA, Meier A, Greco N, Zhang W, Liu W, Lenz RA: Efficacy and safety evaluation of hsd-1 inhibitor abt-384 in alzheimer's disease. *Alzheimer's and Dementia* 2014;10:S364-S373.
- 33 Peng DT, Xu XH, Wang LN: Efficiency and safety assessment of donepezil for treating mild and moderate alzheimer disease. *Chinese Journal of Clinical Rehabilitation* 2005;9:170-172.
- 34 Rogers SL, Doody RS, Mohs RC, Friedhoff LT, Donepezil Study G: Donepezil improves cognition and global function in alzheimer disease - a 15-week, double-blind, placebo-controlled study. *Arch Intern Med* 1998;158:1021-1031.
- 35 Howard RJ, Juszczak E, Ballard CG, Bentham P, Brown RG, Bullock R, Burns AS, Holmes C, Jacoby R, Johnson T, Knapp M, Lindesay J, O'Brien JT, Wilcock G, Katona C, Jones RW, DeCesare J, Rodger M, Grp CAT: Donepezil for the treatment of agitation in alzheimer's disease. *N Engl J Med* 2007;357:1382-1392.
- 36 Moraes W, Poyares D, Sukys-Claudino L, Guilleminault C, Tufik S: Donepezil improves obstructive sleep apnea in alzheimer disease - a double-blind, placebo-controlled study. *Chest* 2008;133:677-683.

- 37 Sole-Padulles C, Bartres-Faz D, Llado A, Bosch B, Pena-Gomez C, Castellvi M, Rami L, Bargallo N, Sanchez-Valle R, Luis Molinuevo J: Donepezil treatment stabilizes functional connectivity during resting state and brain activity during memory encoding in alzheimer's disease. *Journal of Clinical Psychopharmacology* 2013;33:199-205.
- 38 Haig GM, Pritchett Y, Meier A, Othman AA, Hall C, Gault LM, Lenz RA: A randomized study of h3 antagonist abt-288 in mild-to-moderate alzheimer's dementia. *Journal of Alzheimer's Disease* 2014;42:959-971.
- 39 Black SE, Doody R, Li H, McRae T, Jambor KM, Xu Y, Sun Y, Perdomo CA, Richardson S: Donepezil preserves cognition and global function in patients with severe alzheimer disease. *Neurology* 2007;69:459-469.
- 40 Burns A, Rossor M, Hecker J, Gauthier S, Petit H, Moller HJ, Rogers SL, Friedhoff LT, Int Donepezil Study G: The effects of donepezil in alzheimer's disease - results from a multinational trial. *Dementia and Geriatric Cognitive Disorders* 1999;10:237-244.
- 41 Feldman H, Gauthier S, Hecker J, Vellas B, Subbiah P, Whalen E, Grp DMSI: A 24-week, randomized, double-blind study of donepezil in moderate to severe alzheimer's disease. *Neurology* 2001;57:613-620.
- 42 Gold M, Alderton C, Zvartau-Hind M, Egginton S, Saunders AM, Irizarry M, Craft S, Landreth G, Linnamagi U, Sawchak S: Rosiglitazone monotherapy in mild-to-moderate alzheimer's disease: Results from a randomized, double-blind, placebo-controlled phase iii study. *Dementia and Geriatric Cognitive Disorders* 2010;30:131-146.
- 43 Homma A, Takeda M, Imai Y, Udaka F, Hasegawa K, Kameyama M, Nishimura T, Grp ES: Clinical efficacy and safety of donepezil on cognitive and global function in patients with alzheimer's disease - a 24-week, multicenter, double-blind, placebo-controlled study in japan. *Dementia and Geriatric Cognitive Disorders* 2000;11:299-313.
- 44 Jia J, Wei C, Jia L, Tang Y, Liang J, Zhou A, Li F, Shi L, Doody RS: Efficacy and safety of donepezil in chinese patients with severe alzheimer's disease: A randomized controlled trial. *Journal of Alzheimer's Disease* 2017;56:1495-1504.
- 45 Maher-Edwards G, Dixon R, Hunter J, Gold M, Hopton G, Jacobs G, Hunter J, Williams P: Sb-742457 and donepezil in alzheimer disease: A randomized, placebo-controlled study. *International Journal of Geriatric Psychiatry* 2011;26:536-544.
- 46 Mazza M, Capuano A, Bria P, Mazza S: Ginkgo biloba and donepezil: A comparison in the treatment of alzheimer's dementia in a randomized placebo-controlled double-blind study. *Eur J Neurol* 2006;13:981-985.
- 47 Gault LM, Lenz RA, Ritchie CW, Meier A, Othman AA, Tang Q, Berry S, Pritchett Y, Robieson WZ: Abt-126 monotherapy in mild-to-moderate alzheimer's dementia: Randomized double-blind, placebo and active controlled adaptive trial and open-label extension. *Alzheimers Res Ther* 2016;8
- 48 Rogers SL, Farlow MR, Doody RS, Mohs R, Friedhoff LT, Albala B, Baumel B, Booker G, Dexter J, Farmer M, Feighner JP, Ferris S, Gordon B, Gorman DG, Hanna G, Harrell LE, Hubbard R, Kennedy J, McCarthy J, Scharre DW, Schaerf F, Schneider L, Seltzer B, Siegal A, Stark SR, Strauss A, Walshe TM, Donepezil Study G: A 24-week, double-blind, placebo-controlled trial of donepezil in patients with alzheimer's disease. *Neurology* 1998;50:136-145.
- 49 Seltzer B, Zolnouni P, Nunez M, Goldman R, Kumar D, Ieni J, Richardson S, Grp DS: Efficacy of donepezil in early-stage alzheimer disease - a randomized placebo-controlled trial. *Archives of Neurology* 2004;61:1852-1856.
- 50 Tune L, Tiseo PJ, Ieni J, Perdomo C, Pratt RD, Votaw JR, Jewart RD, Hoffman JM: Donepezil hcl (e2020) maintains functional brain activity in patients with alzheimer disease - results of a 24-week, double-blind, placebo-controlled study. *American Journal of Geriatric Psychiatry* 2003;11:169-177.

- 51 Maher-Edwards G, Watson C, Ascher J, Barnett C, Boswell D, Davies J, Fernandez M, Kurz A, Zanetti O, Safirstein B, Schronen JP, Zvartau-Hind M, Gold M: Two randomized controlled trials of sb742457 in mild-to-moderate alzheimer's disease. *Alzheimer's and Dementia: Translational Research and Clinical Interventions* 2015;1:23-36.
- 52 dos Santos Moraes WA, Poyares DR, Guilleminault C, Ramos LR, Ferreira Bertolucci PH, Tufik S: The effect of donepezil on sleep and rem sleep eeg in patients with alzheimer disease: A double-blind placebo-controlled study. *Sleep* 2006;29:199-205.
- 53 Winblad B, Kilander L, Eriksson S, Minthon L, Batsman S, Wetterholm AL, Jarisson-Blixt C, Hoglund A, Severe Alzheimers Dis Study G: Donepezil in patients with severe alzheimer's disease: Double-blind, parallel-group, placebo-controlled study. *Lancet* 2006;367:1057-1065.
- 54 Winblad B, Engedal K, Soininen H, Verhey F, Waldemar G, Wimo A, Wetterholm AL, Zhang R, Haglund A, Subbiah P, Grp DNS: A 1-year, randomized, placebo-controlled study of donepezil in patients with mild to moderate ad. *Neurology* 2001;57:489-495.
- 55 Mohs RC, Doody RS, Morris JC, Ieni JR, Rogers SL, Perdomo CA, Pratt RD, Grp S: A 1-year, placebo-controlled preservation of function survival study of donepezil in ad patients. *Neurology* 2001;57:481-488.
- 56 Bentham P, Gray R, Raftery J, Hills R, Sellwood E, Courtney C, Farrell D, Hardyman W, Crome P, Edwards S, Lendon C, Lynch L, Grp ADC: Long-term donepezil treatment in 565 patients with alzheimer's disease (ad2000): Randomised double-blind trial. *Lancet* 2004;363:2105-2115.
- 57 Tariot PN, Cummings JL, Katz IR, Mintzer J, Perdomo CA, Schwam EM, Whalen E: A randomized, double-blind, placebo-controlled study of the efficacy and safety of donepezil in patients with alzheimer's disease in the nursing home setting. *Journal of the American Geriatrics Society* 2001;49:1590-1599.
- 58 Black S, Roman GC, Geldmacher DS, Salloway S, Hecker J, Burns A, Perdomo C, Kumar D, Pratt R, Donepezil 307 Vascular Dementia S: Efficacy and tolerability of donepezil in vascular dementia - positive results of a 24-week, multicenter, international, randomized, placebo-controlled clinical trial. *Stroke* 2003;34:2323-2330.
- 59 Roman GC, Salloway S, Black SE, Royall DR, DeCarli C, Weiner MW, Moline M, Kumar D, Schindler R, Posner H: Randomized, placebo-controlled, clinical trial of donepezil in vascular dementia differential effects by hippocampal size. *Stroke* 2010;41:1213-1221.
- 60 Wilkinson D, Doody R, Helme R, Taubman K, Mintzer J, Kertesz A, Pratt RD, Donepezil 308 Study G: Donepezil in vascular dementia - a randomized, placebo-controlled study. *Neurology* 2003;61:479-486.
- 61 Dichgans M, Markus HS, Salloway S, Verkkoniemi A, Moline M, Wang Q, Posner H, Chabriat HS: Donepezil in patients with subcortical vascular cognitive impairment: A randomised double-blind trial in cadasil. *The Lancet Neurology* 2008;7:310-318.
- 62 Aarsland D, Laake K, Larsen JP, Janvin C: Donepezil for cognitive impairment in parkinson's disease: A randomised controlled study. *Journal of Neurology Neurosurgery and Psychiatry* 2002;72:708-712.
- 63 Ravina B, Putt M, Siderowf A, Farrar JT, Gillespie M, Crawley A, Fernandez HH, Trieschmann MM, Reichwein S, Simuni T: Donepezil for dementia in parkinson's disease: A randomised, double blind, placebo controlled, crossover study. *Journal of Neurology Neurosurgery and Psychiatry* 2005;76:934-939.
- 64 Leroi I, Brandt J, Reich SG, Lyketsos CG, Grill S, Thompson R, Marsh L: Randomized placebo-controlled trial of donepezil in cognitive impairment in parkinson's disease. *International Journal of Geriatric Psychiatry* 2004;19:1-8.

- 65 Dubois B, Tolosa E, Katzenschlager R, Emre M, Lees AJ, Schumann G, Pourcher E, Gray J, Thomas G, Swartz J, Hsu T, Moline ML: Donepezil in parkinson's disease dementia: A randomized, double-blind efficacy and safety study. *Movement Disorders* 2012;27:1230-1238.
- 66 Ikeda M, Mori E, Matsuo K, Nakagawa M, Kosaka K: Donepezil for dementia with lewy bodies: A randomized, placebo-controlled, confirmatory phase iii trial. *Alzheimers Res Ther* 2015;7
- 67 Mori E, Ikeda M, Kosaka K, Donepezil DLBSI: Donepezil for dementia with lewy bodies: A randomized, placebo-controlled trial. *Annals of Neurology* 2012;72:41-52.
- 68 Wilkinson D, Murray J, Grp GR: Galantamine: A randomized, double-blind, dose comparison in patients with alzheimer's disease. *International Journal of Geriatric Psychiatry* 2001;16:852-857.
- 69 Kadir A, Darreh-Shori T, Almkvist O, Wall A, Grut M, Strandberg B, Ringheim A, Eriksson B, Blomquist G, Langstrom B, Nordberg A: Pet imaging of the in vivo brain acetylcholinesterase activity and nicotine binding in galantamine-treated patients with ad. *Neurobiology of Aging* 2008;29:1204-1217.
- 70 Rockwood K, Mintzer J, Truyen L, Wessel T, Wilkinson D: Effects of a flexible galantamine dose in alzheimer's disease: A randomised, controlled trial. *Journal of Neurology Neurosurgery and Psychiatry* 2001;71:589-595.
- 71 Rockwood K, Fay S, Song XW, MacKnight C, Gorman M, Investigators V: Attainment of treatment goals by people with alzheimer's disease receiving galantamine: A randomized controlled trial. *Canadian Medical Association Journal* 2006;174:1099-1105.
- 72 Tariot PN, Solomon PR, Morris JC, Kershaw P, Lilienfeld S, Ding C, Grp GU-S: A 5-month, randomized, placebo-controlled trial of galantamine in ad. *Neurology* 2000;54:2269-2276.
- 73 Brodaty H, Corey-Bloom J, Potocnik FCV, Truyen L, Gold M, Damaraju CRV: Galantamine prolonged-release formulation in the treatment of mild to moderate alzheimer's disease. *Dementia and Geriatric Cognitive Disorders* 2005;20:120-132.
- 74 Raskind MA, Peskind ER, Wessel T, Yuan W, Grp GU-S: Galantamine in ad - a 6-month randomized, placebo-controlled trial with a 6-month extension. *Neurology* 2000;54:2261-2268.
- 75 Wilcock GK, Lilienfeld S, Gaens E, Study GI-: Efficacy and safety of galantamine in patients with mild to moderate alzheimer's disease: Multicentre randomised controlled trial. *British Medical Journal* 2000;321:1445-1449.
- 76 Likitjaroen Y, Meindl T, Frieze U, Wagner M, Buerger K, Hampel H, Teipel SJ: Longitudinal changes of fractional anisotropy in alzheimer's disease patients treated with galantamine: A 12-month randomized, placebo-controlled, double-blinded study. *Eur Arch Psych Clin Neurosci* 2012;262:341-350.
- 77 Hager K, Baseman AS, Nye JS, Brashear HR, Han J, Sano M, Davis B, Richards HM: Effects of galantamine in a 2-year, randomized, placebo-controlled study in alzheimer's disease. *Neuropsychiatric Disease and Treatment* 2014;10:391-401.
- 78 Erkinjuntti T, Kurz A, Gauthier S, Bullock R, Lilienfeld S, Damaraju CV: Efficacy of galantamine in probable vascular dementia and alzheimer's disease combined with cerebrovascular disease: A randomised trial. *Lancet* 2002;359:1283-1290.
- 79 Auchus AP, Brashear HR, Salloway S, Korczyn AD, De Deyn PP, Gassmann-Mayer C, Grp G-I-S: Galantamine treatment of vascular dementia: A randomized trial. *Neurology* 2007;69:448-458.
- 80 Litvinenko IV, Odinak MM, Mogil'naya VI, Emelin AY: Efficacy and safety of galantamine (reminyl) for dementia in patients with parkinson's disease (an open controlled trial). *Neuroscience and behavioral physiology* 2008;38:937-945.
- 81 Koch G, Di Lorenzo F, Bonni S, Giacobbe V, Bozzali M, Caltagirone C, Martorana A: Dopaminergic modulation of cortical plasticity in alzheimer's disease patients. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology* 2014;39:2654-2661.
- 82 Mowla A, Mosavinasab M, Haghshenas H, Borhani-Haghighi A: Does serotonin augmentation have any effect on cognition and activities of daily living in alzheimer's dementia? - a double-blind, placebo-controlled clinical trial. *Journal of Clinical Psychopharmacology* 2007;27:484-487.

- 83 Iranmanesh F, Vakilian A, Gadari F, Sayyadi A, Mehrabian M, Moradi M, Raesy E: Piracetam, rivastigmine and their joint consumption effects on mmse score status in patients with alzheimer's disease. 2013 2012;11:60-63.
- 84 Agid Y, Dubois B, Anand R, Gharabawi G, Int Rivastigmine I: Efficacy and tolerability of rivastigmine in patients with dementia of the alzheimer type. *Curr Ther Res-Clin Exp* 1998;59:837-845.
- 85 Forette F, Anand R, Gharabawi G: A phase ii study in patients with alzheimer's disease to assess the preliminary efficacy and maximum tolerated dose of rivastigmine (exelon (r)). *Eur J Neurol* 1999;6:423-429.
- 86 Winblad B, Cummings J, Andreasen N, Grossberg G, Onofrj M, Sadowsky C, Zechner S, Nagel J, Lane R: A six-month double-blind, randomized, placebo-controlled study of a transdermal patch in alzheimer's disease - rivastigmine patch versus capsule. *International Journal of Geriatric Psychiatry* 2007;22:456-467.
- 87 Rosler M, Anand R, Cicin-Sain A, Gauthier S, Agid Y, Dal-Bianco P, Stahelin HB, Hartman R, Gharabawi M, Grp BES: Efficacy and safety of rivastigmine in patients with alzheimer's disease: International randomised controlled trial. *British Medical Journal* 1999;318:633-638.
- 88 Corey-Bloom J, Anand R, Veach J: A randomized trial evaluating the efficacy and safety of ena 713 (rivastigmine tartrate), a new acetylcholinesterase inhibitor, in patients with mild to moderately severe alzheimer's disease. *International Journal of Geriatric Psychopharmacology* 1998;1:55-65.
- 89 Feldman HH, Lane R: Rivastigmine: A placebo controlled trial of twice daily and three times daily regimens in patients with alzheimer's disease. *Journal of Neurology Neurosurgery and Psychiatry* 2007;78:1056-1063.
- 90 Karaman Y, Erdogan F, Koseoglu E, Turan T, Ersoy AO: A 12-month study of the efficacy of rivastigmine in patients with advanced moderate alzheimer's disease. *Dementia and Geriatric Cognitive Disorders* 2005;19:51-56.
- 91 Ballard C, Sauter M, Scheltens P, He Y, Barkhof F, van Straaten ECW, van der Flier WM, Hsu C, Wu S, Lane R: Efficacy, safety and tolerability of rivastigmine capsules in patients with probable vascular dementia: The vantage study. *Current Medical Research and Opinion* 2008;24:2561-2574.
- 92 Mok V, Wong A, Ho S, Leung T, Lam WWM, Wong KS: Rivastigmine in chinese patients with subcortical vascular dementia. *Neuropsychiatric disease and treatment* 2007;3:943-948.
- 93 Emre M, Aarsland D, Albanese A, Byrne EJ, Deuschl G, De Deyn PP, Durif F, Kulisevsky J, van Laar T, Lees A, Poewe W, Robillard A, Rosa MM, Wolters E, Quarg P, Tekin S, Lane R: Rivastigmine for dementia associated with parkinson's disease. *N Engl J Med* 2004;351:2509-2518.
- 94 Fox C, Crugel M, Maidment I, Auestad BH, Coulton S, Treloar A, Ballard C, Boustani M, Katona C, Livingston G: Efficacy of memantine for agitation in alzheimer's dementia: A randomised double-blind placebo controlled trial. *PLoS One* 2012;7
- 95 Bakchine S, Loft H: Memantine treatment in patients with mild to moderate alzheimer's disease: Results of a randomised, double-blind, placebo-controlled 6-month study. *Journal of Alzheimers Disease* 2008;13:97-107.
- 96 Peskind ER, Potkin SG, Pomara N, Ott BR, Graham SM, Olin JT, McDonald S, Grp M-M-S: Memantine treatment in mild to moderate alzheimer disease: A 24-week randomized, controlled trial. *American Journal of Geriatric Psychiatry* 2006;14:704-715.
- 97 Wang T, Huang Q, Reiman EM, Chen K, Li X, Li G, Lin Z, Li C, Xiao S: Effects of memantine on clinical ratings, fluorodeoxyglucose positron emission tomography measurements, and cerebrospinal fluid assays in patients with moderate to severe alzheimer dementia: A 24-week, randomized, clinical trial. *Journal of Clinical Psychopharmacology* 2013;33:636-642.
- 98 Reisberg B, Doody R, Stoffler A, Schmitt F, Ferris S, Mobius HJ, Memantine Study G: Memantine in moderate-to-severe alzheimer's disease. *N Engl J Med* 2003;348:1333-1341.

- 99 Ashford JW, Adamson M, Beale T, La D, Hernandez B, Noda A, Rosen A, O'Hara R, Fairchild JK, Spielman D, Yesavage JA: Mr spectroscopy for assessment of memantine treatment in mild to moderate alzheimer dementia. *Journal of Alzheimers Disease* 2011;26:331-336.
- 100 Wilkinson D, Fox NC, Barkhof F, Phul R, Lemming O, Scheltens P, Study I: Memantine and brain atrophy in alzheimer's disease: A 1-year randomized controlled trial. *Journal of Alzheimers Disease* 2012;29:459-469.
- 101 Orgogozo JM, Rigaud AS, Stoffler A, Mobius HJ, Forette F: Efficacy and safety of memantine in patients with mild to moderate vascular dementia - a randomized, placebo-controlled trial (mmm 300). *Stroke* 2002;33:1834-1839.
- 102 Wilcock G, Mobius HJ, Stoffler A, Grp MMM: A double-blind, placebo-controlled multicentre study of memantine in mild to moderate vascular dementia (mmm500). *Int Clin Psychopharmacol* 2002;17:297-305.
- 103 Leroi I, Overshott R, Byrne EJ, Daniel E, Burns A: Randomized controlled trial of memantine in dementia associated with parkinson's disease. *Movement Disorders* 2009;24:1217-1221.
- 104 Aarsland D, Ballard C, Walker Z, Bostrom F, Alves G, Kossakowski K, Leroi I, Pozo-Rodriguez F, Minthon L, Londos E: Memantine in patients with parkinson's disease dementia or dementia with lewy bodies: A double-blind, placebo-controlled, multicentre trial. *Lancet Neurol* 2009;8:613-618.
- 105 Boxer AL, Knopman DS, Kaufer DI, Grossman M, Onyike C, Graf-Radford N, Mendez M, Kerwin D, Lerner A, Wu CK, Koestler M, Shapira J, Sullivan K, Klepac K, Lipowski K, Ullah J, Fields S, Kramer JH, Merrilees J, Neuhaus J, Mesulam MM, Miller BL: Memantine in patients with frontotemporal lobar degeneration: A multicentre, randomised, double-blind, placebo-controlled trial. *Lancet Neurol* 2013;12:149-156.
- 106 Vercelletto M, Boutoleau-Bretonniere C, Volteau C, Puel M, Auriacombe S, Sarazin M, Michel BF, Couratier P, Thomas-Anterion C, Verpillat P, Gabelle A, Golfier V, Cerato E, Lacomblez L: Memantine in behavioral variant frontotemporal dementia: Negative results. *Journal of Alzheimers Disease* 2011;23:749-759.

Table 1: Characteristics of included studies (CVD=cerebrovascular disease, CADASIL=Cerebral Autosomal-Dominant Arteriopathy with Subcortical Infarcts and Leukoencephalopathy, PRC=prolonged-release capsule, BID=twice daily, TID=three-times daily)

| Study | Diagnosis | Duration (weeks) | Cognitive measure | Trial arms (n) | Risk of bias |
|--------------------------------|-----------|------------------|-------------------|----------------------------------------------|--------------|
| Donepezil | | | | | |
| Frolich et al., 2011[29] | AD | 12 | MMSE | 5 or 10mg/d (161) Placebo (164) | Unclear |
| Gault et al., 2015[30] | AD | 12 | ADAS-cog | 10mg/d (68) Placebo (68) | Low |
| Gelmacher et al., 2000[31] | AD | 12 | MMSE | Donepezil (6) Placebo (6) | Unclear |
| Marek et al., 2014[32] | AD | 12 | MMSE | 10mg/d (66) Placebo (66) | High |
| Peng et al., 2005[33] | AD | 12 | MMSE | 5mg/d (46) Placebo (43) | High |
| Rogers et al., 1998a[34] | AD | 12 | MMSE | 5mg/d (157) 10mg/d (158) Placebo (153) | High |
| NCT00777608 | AD | 12 | ADAS-cog | 5 or 10mg/d (53) Placebo (53) | High |
| Howard et al., 2007[35] | AD | 12 | MMSE | 10mg/d (128) Placebo (131) | Low |
| Moraes et al., 2008[36] | AD | 13 | ADAS-cog | 5mg/d (11) Placebo (12) | Unclear |
| Sole-Padulles et al., 2013[37] | AD | 13 | MMSE | 10mg/d (8) Placebo (7) | High |
| Haig et al., 2014[38] | AD | 14 | MMSE | 10mg/d (60) Placebo (63) | Low |
| Black et al., 2007[39] | AD | 24 | MMSE | 10mg/d (176) Placebo (167) | High |
| Burns et al., 1999[40] | AD | 24 | ADAS-cog | 5mg/d (271) 10mg/d (273) Placebo (274) | Unclear |
| Feldman et al., 2000[41] | AD | 24 | MMSE | 10mg/d (144) Placebo (146) | Unclear |
| Gold et al., 2010[42] | AD | 24 | ADAS-cog | 10mg/d (84) Placebo (166) | High |
| Homma et al., 2000[43] | AD | 24 | ADAS-cog | 5mg/d (134) Placebo (129) | Unclear |
| Jia et al., 2017[44] | AD | 24 | MMSE | 5mg/d (156) Placebo (156) | Low |
| Maher-Edwards et al., 2011[45] | AD | 24 | ADAS-cog | 10mg/d (67) Placebo (63) | High |
| Mazza et al., 2006[46] | AD | 24 | MMSE | 5mg/d (25) Placebo (26) | High |

| | | | | | |
|------------------------------------|-----------------|----|------------------|----------------------------------------------|---------|
| Gault et al., 2016[47] | AD | 24 | MMSE | 10mg/d (76) Placebo (104) | Unclear |
| Rogers et al., 1998b[48] | AD | 24 | MMSE ADAS-cog | 5mg/d (154) 10mg/d (157) Placebo (162) | High |
| Seltzer et al., 2004[49] | AD | 24 | MMSE ADAS-cog | 10mg/d (96) Placebo (57) | High |
| Tune et al., 2003[50] | AD | 24 | ADAS-cog | 10mg/d (14) Placebo (14) | Unclear |
| Maher-Edwards et al., 2015[51] | AD | 24 | MMSE ADAS-cog | 5 or 10mg/d (152) Placebo (145) | High |
| dos Santos Moraes et al., 2006[52] | AD | 26 | ADAS-cog | 10mg/d (17) Placebo (18) | Low |
| Winblad et al., 2006[53] | AD | 26 | MMSE | 10mg/d (128) Placebo (121) | High |
| Winblad et al., 2001[54] | AD | 52 | MMSE | 10mg/d (142) Placebo (144) | Unclear |
| Mohs et al., 2001[55] | AD | 54 | MMSE | 10mg/d (214) Placebo (217) | High |
| Bentham et al., 2004[56] | AD or AD+VaD | 12 | MMSE | 5mg/d (282) Placebo (283) | High |
| Tariot et al., 2001[57] | AD or AD+CVD | 24 | MMSE | 10mg/d (103) Placebo (105) | High |
| Black et al., 2003[58] | VaD | 24 | MMSE ADAS-cog | 5mg/d (198) 10mg/d (206) Placebo (199) | High |
| Roman et al., 2010[59] | VaD | 24 | MMSE | 5mg/d (648) Placebo (326) | High |
| Wilkinson et al., 2003[60] | VaD | 24 | MMSE | 5mg/d (208) 10mg/d (215) Placebo (193) | High |
| Dichgans et al., 2008[61] | CADASIL | 18 | MMSE | 10mg/d (86) Placebo (82) | Unclear |
| Aarsland et al., 2002[62] | PDD | 10 | MMSE | 5 or 10mg/d (8) Placebo (6) | High |
| Ravina et al., 2005[63] | PDD | 10 | ADAS-cog | 5mg/d (11) Placebo (11) | High |
| Leroi et al., 2004[64] | PDD | 18 | MMSE | 10mg/d (7) Placebo (9) | Unclear |
| Dubois et al., 2012[65] | PDD | 24 | MMSE ADAS-cog | 5mg/d (195) 10mg/d (182) Placebo (173) | High |
| Ikeda et al., 2015[66] | DLB | 12 | MMSE | 5mg/d (46) 10mg/d (47) Placebo (49) | High |
| Mori et al., 2012[67] | DLB | 12 | MMSE | 3mg/d (35) 5mg/d (33) | Low |

| | | | | | |
|--------------------------------|------------------|-----|----------|--------------------------------------------------------------|---------|
| | | | | 10mg/d (37) Placebo (35) | |
| <u>Galantamine</u> | | | | | |
| Wilkinson and Murray, 2001[68] | AD | 12 | ADAS-cog | 18mg/d (88) 24mg/d (56) 36mg/d (54) Placebo (87) | High |
| Kadir et al., 2008[69] | AD | 13 | MMSE | 8-16mg/d (12) Placebo (6) | Unclear |
| Rockwood et al., 2001[70] | AD | 13 | ADAS-cog | 24-32mg/d (261) Placebo (125) | High |
| Rockwood et al., 2006[71] | AD | 16 | ADAS-cog | 16-24mg/d (64) Placebo (66) | Unclear |
| Tariot et al., 2000[72] | AD | 22 | ADAS-cog | 8mg/d (140) 16mg/d (279) 24mg/d (273) Placebo (286) | Unclear |
| Brodaty et al., 2005[73] | AD | 26 | ADAS-cog | 16-24mg/d (237) 16-24mg/d PRC (320) Placebo (324) | High |
| Raskind et al., 2000[74] | AD | 26 | ADAS-cog | 24mg/d (212) 32mg/d (211) Placebo (213) | High |
| Wilcock et al., 2000[75] | AD | 26 | ADAS-cog | 24mg/d (220) 32mg/d (218) Placebo (215) | High |
| Likitjaroen et al., 2011[76] | AD | 26 | MMSE | 16mg/d (14) Placebo (11) | Unclear |
| Hager et al., 2014[77] | AD or AD+CVD | 104 | MMSE | 18-24mg/d (1028) Placebo (1023) | Low |
| Erkinjuntti et al., 2002[78] | VaD or AD+CVD | 26 | ADAS-cog | 24mg/d (396) Placebo (196) | High |
| Auchus et al., 2007[79] | VaD | 26 | ADAS-cog | 24mg/d (397) Placebo (391) | High |
| Litvinenko et al., 2008[80] | PDD | 24 | MMSE | 16mg/d (21) Placebo (20) | High |
| <u>Rivastigmine</u> | | | | | |
| Koch et al., 2014[81] | AD | 4 | MMSE | 4.6mg/d (10) Placebo (10) | Unclear |
| Mowla et al., 2007[82] | AD | 12 | MMSE | 6-12mg/d (41) Placebo (40) | Unclear |
| Iranmanesh et al., 2012[83] | AD | 12 | MMSE | 3mg/d (16) Placebo (16) | Unclear |
| Agid et al., 1998[84] | AD | 13 | MMSE | 4mg/d (136) 6mg/d (133) Placebo (133) | High |
| Forette et al., 1999[85] | AD | 18 | ADAS-cog | 12mg/d BID (45) 12mg/d TID (45) | High |

| | | | | | |
|------------------------------|---------|----|------------------|------------------------------------------------------------------------------------------------------|---------|
| Winblad et al., 2007[86] | AD | 24 | MMSE | Placebo (24) 12mg/d capsule (297) 9.5mg/d patch (293) 17.4mg/d patch (303) Placebo (302) | High |
| NCT00423085 | AD | 24 | MMSE | 9mg/d patch (284) 18mg/d patch (287) Placebo (288) | High |
| Rosler et al., 1999[87] | AD | 26 | MMSE | 1-4mg/d (243) 6-12 mg/d (243) Placebo (239) | High |
| Corey-Bloom et al., 1998[88] | AD | 26 | MMSE ADAS-cog | 1-4mg/d (233) 6-12 mg/d (231) Placebo (235) | High |
| Feldman and Lane, 2007[89] | AD | 26 | MMSE ADAS-cog | 2-12mg/d BID (229) 2-12mg/d TID (227) Placebo (222) | Unclear |
| Karaman et al., 2005[90] | AD | 52 | MMSE | 12mg/d (24) Placebo (20) | High |
| Ballard et al., 2008[91] | VaD | 24 | MMSE | 3-12mg/d (365) Placebo (345) | High |
| Mok et al., 2007[92] | VaD | 26 | MMSE | 6mg/d (20) Placebo (20) | Unclear |
| Emre et al., 2004[93] | PDD | 24 | MMSE | 3-12mg/d (362) Placebo (179) | High |
| Memantine | | | | | |
| Fox et al., 2012[94] | AD | 12 | MMSE | 20mg/d (74) Placebo (79) | Low |
| Bakchine and Loft, 2007[95] | AD | 24 | ADAS-cog | 20mg/d (318) Placebo (152) | Low |
| Peskind et al., 2006[96] | AD | 24 | ADAS-cog | 20mg/d (201) Placebo (202) | Low |
| Wang et al., 2013[97] | AD | 24 | MMSE | 20mg/d (13) Placebo (13) | Unclear |
| Reisberg et al., 2003[98] | AD | 28 | MMSE | 20mg/d (126) Placebo (126) | High |
| Ashford et al., 2011[99] | AD | 52 | ADAS-cog | 20mg/d (7) Placebo (6) | High |
| Wilkinson et al., 2012[100] | AD | 52 | MMSE | 20mg/d (134) Placebo (144) | Low |
| Orgogozo et al., 2002[101] | VaD | 28 | MMSE | 20mg/d (165) Placebo (156) | High |
| Wilcock et al., 2002[102] | VaD | 28 | MMSE ADAS-cog | 20mg/d (295) Placebo (284) | Low |
| Leroi et al., 2009[103] | PDD | 16 | MMSE | 20mg/d (11) Placebo (14) | High |
| Aarsland et al., 2009[104] | PDD/DLB | 24 | MMSE | 20mg/d (35) Placebo (40) | Low |

| | | | | | |
|----------------------------------|-----|----|------|-----------------------------|------|
| Boxer et al., 2013[105] | FTD | 26 | MMSE | 20mg/d (39) Placebo (42) | Low |
| Vercelletto et al., 2011[106] | FTD | 52 | MMSE | 20mg/d (26) Placebo (26) | High |

Table 2: Meta-regressions of effects at 3 months. Coefficients, associated p-values and omnibus test of parameters p-value provided. *=significant at 5% level. **=significant at 1% level. ref=reference category.

| Factor | Levels | Number of trials | Coefficient (p-value) | Omnibus test p-value |
|---------------------|--------------|------------------|-----------------------|----------------------|
| Translation to MMSE | MMSE | 28 | ref | 0.007** |
| | ADAS-cog | 32 | -0.471 (0.007) | |
| Risk of bias rating | Low | 8 | ref | 0.521 |
| | Unclear | 13 | -0.371 (0.307) | |
| | High | 39 | -0.346 (0.269) | |
| Medication | Donepezil | 37 | ref | 0.864 |
| | Galantamine | 17 | 0.010 (0.961) | |
| | Rivastigmine | 6 | -0.153 (0.612) | |
| Diagnosis | AD | 46 | ref | 0.009** |
| | VaD | 6 | -0.211 (0.373) | |
| | PDD/DLB | 8 | 0.806 (0.005) | |
| Baseline MMSE | NA | 55 | -0.069 (0.092) | 0.092 |
| Date | Pre 2000 | 26 | ref | 0.703 |
| | 2000 onwards | 34 | 0.068 (0.703) | |

Table 3: Meta-regressions of effects at 6 months. Coefficients, associated p-values and omnibus test of parameters p-value provided. *=significant at 5% level. **=significant at 1% level. ref=reference category.

| Factor | Levels | Number of trials | Coefficient (p-value) | Omnibus test p-value |
|---------------------|--------------|------------------|-----------------------|----------------------|
| Translation to MMSE | MMSE | 35 | ref | 0.540 |
| | ADAS-cog | 17 | 0.117 (0.540) | |
| Risk of bias rating | Low | 3 | ref | 0.735 |
| | Unclear | 9 | 0.269 (0.579) | |
| | High | 40 | 0.329 (0.443) | |
| Medication | Donepezil | 27 | ref | 0.033* |
| | Galantamine | 11 | 0.320 (0.139) | |
| | Rivastigmine | 14 | -0.370 (0.133) | |
| Diagnosis | AD | 39 | ref | |

| | | | | |
|---------------|--------------------------|----------|---------------------------------|--------|
| | VaD PDD/DLB | 9 4 | -0.134 (0.139) 0.970 (0.001) | 0.007* |
| Baseline MMSE | NA | 52 | -0.005 (0.869) | 0.869 |
| Date | Pre 2000 2000 onwards | 17 35 | ref -0.141 (0.456) | 0.456 |




Figure 1: Flow diagram of trials identified for inclusion in this review through two-tier search strategy.





Figure 2: Forest plot showing treatment effects from individual trials and meta-analysis results for AChEIs at 3 months after treatment initiation






Figure 4: Forest plot showing treatment effects from individual trials and meta-analysis results for AChEIs at 6 months after treatment initiation




Figure 5: Forest plot showing treatment effects from individual trials and meta-analysis results for AChEIs at 12 months after treatment initiation




Figure 6: Forest plots showing treatment effects from individual trials and meta-analysis results for memantine at 3, 6 and 12 months after treatment initiation.

Supplementary Material: Search terms

1. "acetylcholinesterase inhibitor*" OR "cholinesterase inhibitor"
2. donepezil OR arciept* OR E2020
3. galantamine OR galanthamine OR reminyl* OR nivalin* OR razadyne* OR lycoremine
4. rivastigmine OR exelon* OR "SDZ ENA 713"
5. memantine OR namenda* OR ebixa* OR axura* OR memox OR akatinol OR abixa OR "D-145"
OR DMAA OR "DRG-0267"
6. Alzheimer* OR AD
7. "vascular dement*" OR VaD
8. DLB OR LBD OR "lewy* bod"
9. PDD OR "Parkinson* disease dement"
10. Dementia
11. randomi?ed OR trial
12. "meta-analysis" OR "systematic review"

These search terms were adapted to the syntax of the databases used and combined in order to carry out the searches as follows

- Reviews: (1 OR 2 OR 3 OR 4 OR 5) AND (6 OR 7 OR 8 OR 9 OR 10) AND 11 AND 12
- Additional AD searches: (1 OR 2 OR 3 OR 4 OR 5) AND 6 AND 11 AND 12, limit 2013-current
- Additional VaD searches:
 - (1 OR 3 OR 4) AND 7 AND 11 AND 12, limit 2013-current
 - (2 OR 5) AND 7 AND 11 AND 12, limit 2007-current
- Additional PDD/DLB searches:
 - (1 OR 2 OR 3 OR 4) AND (8 OR 9) AND 11 AND 12, limit 2011-current
 - 5 AND (8 OR 9) AND 11 AND 12